VOL. 592, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY 21 FEBRUARY, 1992

CONTENTS

15TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, BASLE, JUNE 3–7, 1991, PART I

Foreword
by F. Erni 1

INTRODUCTION

Separation techniques for biotechnology in the 1990s
by M. Roman and P. R. Brown (Kingston, RI, USA) 3

COLUMN LIQUID CHROMATOGRAPHY

Theory

Increasing data precision in reversed-phase liquid chromatography leads to new information level
by U. Dreyer, H. Melzer and H. J. Möckel (Berlin, Germany) 13

Peak purity, yield and throughput in preparative liquid chromatography with self-displacement
by V. R. Meyer (Berne, Switzerland) 17

Assessment of chromatographic peak purity by means of multi-wavelength detection and correlation-based algorithms
by J. B. Castledine and A. F. Fell (Bradford, UK) and R. Modin and B. Sellberg (Uppsala, Sweden) 27

Theory: optimization

Optimization of the separation of (6*R*)- and (6*S*)-leucovorin and evaluation of the robustness of the optimum
by C. Vandenbosch, C. Vannecke and D. L. Massart (Brussels, Belgium) 37

Use of solvent selectivity optimization procedures for high-performance liquid chromatographic method development
by S. D. Patterson (Sunbury-on-Thames, UK) 43

CRISEBOOK, a Hypermedia version of an expert system for the selection of optimization criteria in high-performance liquid chromatography
by B. Bourguignon, P. Vankeerberghen and D. L. Massart (Brussels, Belgium) 51

Theory: structure–mobility correlations

Chromatographic separation and molecular modelling of triazines with respect to their inhibition of the growth of L1210/R71 cells
by K. Valkó and P. Slégel (Budapest, Hungary) 59

Correlations between high-performance liquid chromatographic retention, X-ray structural and ^{13}C NMR spectroscopic data of flavonoid compounds
by M. C. Pietrogrande, P. Reschiglian and F. Dondi (Ferrara, Italy), Y. D. Kahie (Mogadishu, Somalia) and V. Bertolasi (Ferrara, Italy) 65

Retention behaviour of some ring-substituted phenol derivatives on a porous graphitized carbon column
by E. Forgács, T. Cserháti and K. Valkó (Budapest, Hungary) 75

Materials and methods: sorbents

Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–high-pH buffer eluent. Changes in selectivity with the age of the stationary phase
by R. M. Smith and J. P. Westlake (Loughborough, UK) and R. Gill and M. D. Osselton (Reading, UK) 85

New, stable polyamine-bonded polymer gel column
by N. Hirata, Y. Tamura, M. Kasai, Y. Yanagihara and K. Noguchi (Kawasaki-shi, Japan) 93

Application of a pellicular anion-exchange resin to the separation of inorganic and organic anions by single-column anion exchange
by R. C. Ludwig (Bellafonte, PA, USA) 101

Experimental evaluation of sorbent bed homogeneity in high-performance liquid chromatographic columns
by T. Macko and D. Berek (Bratislava, Czechoslovakia) 109

Investigation of solid surfaces by high-performance liquid chromatography
by A. E. Chalykh, L. N. Kolomiets, O. G. Larionov and N. I. Vinogradova (Moscow, USSR) 121

Materials and methods: eluents

Effect of temperature and mobile phase on the retention of retinoates in reversed-phase liquid chromatography
by M. Salo, H. Vuorela and J. Halmekoski (Helsinki, Finland) 127

Application of the gradient elution technique. Demonstration with a special test mixture and the DryLab G/plus method development software
by R. Däppen (Basle, Switzerland) and I. Molnar (Berlin, Germany) 133

Gradient elution with shorter equilibration times in reversed-phase ion-pair chromatography
by M. Patthy (Budapest, Hungary) 143

Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition
by R. M. Lopes Marques and P. J. Schoenmakers (Eindhoven, Netherlands) 157

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. I. Accuracy of a theory-based model
by J. A. Lewis, D. C. Lommen, W. D. Raddatz, J. W. Dolan and L. R. Snyder (Walnut Creek, CA, USA) and I. Molnar (Berlin, Germany) 183

Computer simulation for the prediction of separation as a function of pH for reversed-phase high-performance liquid chromatography. II. Resolution as a function of simultaneous change in pH and solvent strength
by J. W. Lewis, J. W. Dolan and L. R. Snyder (Walnut Creek, CA, USA) and I. Molnar (Berlin, Germany) 197

Materials and methods: detectors

Nitrogen-specific liquid chromatography detector based on chemiluminescence. Application to the analysis of ammonium nitrogen in waste water
by E. M. Fujinari (Houston, TX, USA) and L. O. Courthaudon (Düsseldorf, Germany) 209

Practical on-line determination of biopolymer molecular weights by high-performance liquid chromatography with classical light-scattering detection
by G. Dollinger, B. Cunico, M. Kunitani, D. Johnson and R. Jones (Emeryville, CA, USA) 215

Constant-potential amperometric detection of carbohydrates at metal electrodes in high-performance anion-exchange chromatography
by T. Ueda, R. Mitchell and F. Kitamura (Lawrence, KS, USA) and A. Nakamoto (Kyoto, Japan) 229

Materials and methods: methods

High-temperature open-tubular capillary column liquid chromatography
by G. Liu, N. M. Djordjevic and F. Erni (Basle, Switzerland) 239

Automated high-performance liquid chromatographic and size-exclusion chromatographic sample preparation by means of a robotic workstation
by A. Bruns, H. Waldhoff, A. Wilsch-Irrgang and W. Winkle (Düsseldorf, Germany) 249

Applications: environmental

Rapid and sensitive determination of urinary 2,5-hexanedione by reversed-phase high-performance liquid chromatography
by M. P. Colombini (Pisa, Italy), P. Carrai (Lucca, Italy) and R. Fuoco and C. Abete (Pisa, Italy) 255

Trace analysis of phthalocyanine pigments by high-performance liquid chromatography
by C.-H. Fischer (Berlin, Germany) 261

Automated determination of aflatoxin B_1 in cattle feed by two-column solid-phase extraction with on-line high-performance liquid chromatography
by J. A. van Rhijn (Wageningen, Netherlands), J. Viveen (Nieuwegein, Netherlands) and L. G. M. Th. Tuinstra (Wageningen, Netherlands) 265

Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products
by G. A. Gross and A. Grüter (Lausanne, Switzerland) 271

Rapid high-performance liquid chromatographic determination of urinary N-(1-methylethyl)-N′-phenyl-1,4-benzenediamine in workers exposed to aromatic amines
by S. Pellegrino and M. Petrarulo (Turin, Italy) and E. Testa and A. Nicolotti (Chieri, Italy) 279

Determination of National Survey of Pesticides analytes in groundwater by liquid chromatography with postcolumn reaction detection
by C. J. Miles (Honolulu, HI, USA) 283

Applications: chiral

Direct separation of enantiomers by high-performance liquid chromatography on a new chiral ligand-exchange phase
by N. Ôi, H. Kitahara and R. Kira (Osaka, Japan) 291

Chiral high-performance liquid chromatographic separations on an α_1-acid glycoprotein column. II. Separation of the diastereomeric and enantiomeric analogues of vinpocetine (Cavinton)
by B. Herényi and S. Görög (Budapest, Hungary) 297

Retention and enantioselectivity of racemic solutes on a modified ovomucoid-bonded column. I. Cross-linking with glutaraldehyde
by J. Haginaka, C. Seyama and H. Yasuda (Nishinomiya, Japan) and H. Fujima and H. Wada (Kyoto, Japan) . . . 301

Investigations of the influence of silanol groups on the separation of enantiomers by liquid and supercritical fluid chromatography
by R. Brügger and H. Arm (Berne, Switzerland) 309

Determination of the enantiomers of fenoldopam in human plasma by reversed-phase high-performance liquid chromatography after chiral derivatization
by V. K. Boppana, L. Geschwindt, M. J. Cyronak and G. Rhodes (King of Prussia, PA, USA) 317

By-products in the derivatization of amines with the chiral reagent 2,3,4,6-tetra-O-acetyl-β-D-glucopyranosyl isothiocyanate and their elimination
by M. Ahnoff, K. Balmér and Y. Lindman (Mölndal, Sweden) 323

Reversed retention order and other stereoselective effects in the separation of amino alcohols on Chiralcel OD
by K. Balmér, P.-O. Lagerström and B.-A. Persson (Mölndal, Sweden) and G. Schill (Uppsala, Sweden) 331

(2*S*,3*S*)-Dicyclohexyl tartrate as mobile phase additive for the determination of the enantiomeric purity of (*S*)-atropine in tablets
by E. Heldin, N. H. Huynh and C. Pettersson (Uppsala, Sweden) 339

High-performance ligand-exchange chromatography of some amino acids containing two chiral centres
by S. V. Galushko, I. P. Shishkina and V. A. Soloshonok (Kiev, Ukraine) 345

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*).